arXiv:hep-ph/9706380v1 17 Jun 1997

Determination of $|V_{td}/V_{ts}|$ within the Standard Model

C. S. Kim
Department of Physics, Yonsei University,
Seoul 120-749, Korea

T. Morozumi [a]
Department of Physics, Hiroshima University,
1-3-1 Kagamiyama, Higashi Hiroshima, Japan 739

A. I. Sanda
Department of Physics, Nagoya University
Chikusa-ku, Nagoya, Japan 464-01

We propose a new method to extract the value of $|V_{td}/V_{ts}|$ from the ratio of the decay distributions $\frac{dBr}{ds}[B \to X_d l^+ l^-]$ and $\frac{dBr}{ds}[B \to X_s l^+ l^-\ ]$.

1 A new method to extract V_{td}/V_{ts}

The determination of the elements of CKM matrix is one of the most important issues in the quark flavor physics. In this talk, we study specifically V_{td}/V_{ts}. In the Standard Model with the unitarity of CKM matrix, V_{ts} is approximated by $-V_{cb}$ which is directly measured by semileptonic B decays. The unknown element $|V_{td}|$ can be extracted through $B_d - \bar{B}_d$ mixing. However, in $B_d - \bar{B}_d$ mixing the uncertainty of the hadronic matrix elements prevents us to extract the element of CKM with good accuracy. In this letter, we propose another method to determine $|V_{td}/V_{ts}|$ with the decay distributions $\frac{dB}{ds}[B \to X_s l^+ l^-\ ]$ and $\frac{dB}{ds}[B \to X_d l^+ l^-]$, where s is invariant mass-squared of $l^+ l^-$. The idea stems from the fact that in the decays $B \to X_q l^+ l^-$ $(q = d, s)$, the short distance (SD) contribution comes from the top quark loop diagrams and the long distance (LD) contribution comes from the decay chains due to charmonium states. The former amplitude is proportional to ${V_{tq}}^* V_{tb}$ and the latter proportional to ${V_{cq}}^* V_{cb}$. If the invariant mass squared of $l^+ l^-$ is away from the peaks of the charmonium resonances (J/ψ and ψ'), the SD contribution is dominant, while on the peaks of the resonances the LD contribution is dominant[1].

In SU(3) limit $m_d = m_s$, the ratio of the distributions

$$\frac{dR}{ds} \equiv \tfrac{dB}{ds}[B \to X_d l^+ l^-] / \tfrac{dB}{ds}[B \to X_s l^+ l^-\ ] \tag{1}$$

---

[a] Talk is presented by T. Morozumi at BCONF97.

becomes $|\frac{V_{td}}{V_{ts}}|^2$, if s is away from the peak of resonance J/ψ. On the peak, the ratio becomes $|\frac{V_{cd}}{V_{cs}}|^2 = \lambda^2$, where λ is Cabibbo angle, $sin\theta_c$. In the intermediate region, there is characteristic interference between the LD contribution and the SD contribution, which requires the detailed study of the distributions. To summarize our prediction in the SU(3) limit:

$$\begin{aligned} \frac{1}{\lambda^2}\frac{dR(s)}{ds} &\to (1-\rho)^2+\eta^2 \quad (s \to 1\mathrm{GeV}^2), \\ &\to \qquad 1 \qquad\quad (s \to m_{J/\psi}{}^2). \end{aligned} \tag{2}$$

In the following, we present the explicit formulae for the distribution as well as the numerical results for the ratio.

2 The differential decay rates and the ratio

Following the notation of Ref. [2], the amplitude of $b \to q l^+ l^-$ $(q = d, s)$ can be written as[3]

$$\begin{aligned} \mathcal{M}(b \to q l^+ l^-\ ) &= \frac{G_F \alpha}{\sqrt{2}\pi} V_{tq}^* V_{tb} \left[\left(C_{9q}^{\mathrm{eff}} - C_{10} \right) (\bar{q}\,\gamma_\mu\, L\, b) \left(\bar{l}\,\gamma^\mu\, L\, l \right) \right. \\ &\quad + \left(C_{9q}^{\mathrm{eff}} + C_{10} \right) (\bar{q}\,\gamma_\mu\, L\, b) \left(\bar{l}\,\gamma^\mu\, R\, l \right) \\ &\quad \left. -2 C_7^{\mathrm{eff}} \left(\bar{q}\, i\, \sigma_{\mu\nu} \frac{q^\nu}{q^2} (m_q L + m_b R)\, b \right) \left(\bar{l}\,\gamma^\mu\, l \right) \right] \end{aligned} \tag{3}$$

where C_{9q}^{eff} is given by,

$$C_{9q}^{\mathrm{eff}}(\hat{s}) \equiv C_9 \left\{ 1 + \frac{\alpha_s(\mu)}{\pi} \omega(\hat{s}) \right\} + Y_{SD}{}^q(\hat{s}) + Y_{LD}{}^q(\hat{s}), \tag{4}$$

and q^ν is the sum of the four momentum of $l^+ l^-$, $s = q^2$ and $\hat{s} = s/m_b{}^2$. The function $Y_{SD}{}^q(\hat{s})$ is the one-loop matrix element of O_9, and $Y_{LD}{}^q(\hat{s})$ is the LD contributions due to the vector mesons J/ψ, ψ' and higher resonances. The function $\omega(\hat{s})$ represents the $O(\alpha_s)$ correction from the one-gluon exchange in the matrix element of O_9. The function $Y_{SD}{}^q$ can be found in the literature[3], which is written as

$$\begin{aligned} Y_{LD}{}^q(\hat{s}) &= \frac{3}{\alpha^2} \kappa \left(-\frac{V_{cq}^* V_{cb}}{V_{tq}^* V_{tb}} C^{(0)} - \frac{V_{uq}^* V_{ub}}{V_{tq}^* V_{tb}} (3\, C_3 + C_4 + 3\, C_5 + C_6) \right) \\ &\quad \times \sum_{V_i = \psi(1s), \ldots, \psi(6s)} \frac{\pi\, \Gamma(V_i \to l^+ l^-)\, M_{V_i}}{M_{V_i}{}^2 - \hat{s}\, m_b{}^2 - i M_{V_i} \Gamma_{V_i}}. \end{aligned} \tag{5}$$

In Eq. (5), $C^{(0)} \simeq 3C_1^{(0)}+C_2^{(0)} \simeq 0.38(0.12)$ while $3C_3^{(0)}+C_4^{(0)}+3\,C_5^{(0)}+C_6^{(0)} \simeq -0.0014(-0.0035)$ for $\mu = 5GeV(2.5GeV)$. Therefore the second term in Y_{LD} can be neglected. The parameter κ is introduced so that the correct value for the branching fraction of the decay chain $B \to X_q J/\psi \to X_q l^+ l^-$ is reproduced. We choose $\kappa C^{(0)} = 0.88$. The differential decay rate for the inclusive semileptonic decay has been calculated for $B \to X_s l^+ l^-$ including the ${\Lambda_{QCD}}^2/{m_b}^2$ power corrections. The result can be translated into $B \to X_q l^+ l^-$.

$$
\begin{aligned}
&\frac{\mathrm{d}\mathcal{B}}{\mathrm{d}\hat{s}} = 2\frac{|V_{tq}^* V_{tb}|^2}{|V_{cb}|^2}\,\mathcal{B}_0 \left\{ \left[\frac{2}{3}\hat{u}(\hat{s},\hat{m}_q)((1-\hat{m}_q^2)^2+\hat{s}(1+\hat{m}_q^2)-2\hat{s}^2)\right.\right.\\
&+\frac{1}{3}(1-4\hat{m}_q^2+6\hat{m}_q^4-4\hat{m}_q^6+\hat{m}_q^8-\hat{s}+\hat{m}_q^2\hat{s}+\hat{m}_q^4\hat{s}-\hat{m}_q^6\hat{s}\\
&-3\hat{s}^2-2\hat{m}_q^2\hat{s}^2-3\hat{m}_q^4\hat{s}^2+5\hat{s}^3+5\hat{m}_q^2\hat{s}^3-2\hat{s}^4)\frac{\hat{\lambda}_1}{\hat{u}(\hat{s},\hat{m}_q)}\\
&+\left(1-8\hat{m}_q^2+18\hat{m}_q^4-16\hat{m}_q^6+5\hat{m}_q^8-\hat{s}-3\hat{m}_q^2\hat{s}+9\hat{m}_q^4\hat{s}-5\hat{m}_q^6\hat{s}-15\hat{s}^2\right.\\
&\left.\left.-18\hat{m}_q^2\hat{s}^2-15\hat{m}_q^4\hat{s}^2+25\hat{s}^3+25\hat{m}_q^2\hat{s}^3-10\hat{s}^4\right)\frac{\hat{\lambda}_2}{\hat{u}(\hat{s},\hat{m}_q)}\right]\left(|C_9^{\mathrm{eff}}|^2+|C_{10}|^2\right)\\
&+\left[\frac{8}{3}\hat{u}(\hat{s},\hat{m}_q)(2(1+\hat{m}_q^2)(1-\hat{m}_q^2)^2-(1+14\hat{m}_q^2+\hat{m}_q^4)\hat{s}-(1+\hat{m}_q^2)\hat{s}^2)\right.\\
&+\frac{4}{3}(2-6\hat{m}_q^2+4\hat{m}_q^4+4\hat{m}_q^6-6\hat{m}_q^8+2\hat{m}_q^{10}\\
&-5\hat{s}-12\hat{m}_q^2\hat{s}+34\hat{m}_q^4\hat{s}-12\hat{m}_q^6\hat{s}-5\hat{m}_q^8\hat{s}+3\hat{s}^2\\
&+29\hat{m}_q^2\hat{s}^2+29\hat{m}_q^4\hat{s}^2+3\hat{m}_q^6\hat{s}^2+\hat{s}^3-10\hat{m}_q^2\hat{s}^3+\hat{m}_q^4\hat{s}^3-\hat{s}^4-\hat{m}_q^2\hat{s}^4)\frac{\hat{\lambda}_1}{\hat{u}(\hat{s},\hat{m}_q)}\\
&+4\left(-6+2\hat{m}_q^2+20\hat{m}_q^4-12\hat{m}_q^6-14\hat{m}_q^8+10\hat{m}_q^{10}+3\hat{s}+16\hat{m}_q^2\hat{s}+62\hat{m}_q^4\hat{s}\right.\\
&-56\hat{m}_q^6\hat{s}-25\hat{m}_q^8\hat{s}+3\hat{s}^2+73\hat{m}_q^2\hat{s}^2+101\hat{m}_q^4\hat{s}^2+15\hat{m}_q^6\hat{s}^2+5\hat{s}^3-26\hat{m}_q^2\hat{s}^3+\\
&\left.\left.5\hat{m}_q^4\hat{s}^3-5\hat{s}^4-5\hat{m}_q^2\hat{s}^4\right)\frac{\hat{\lambda}_2}{\hat{u}(\hat{s},\hat{m}_q)}\right]\frac{|C_7^{\mathrm{eff}}|^2}{\hat{s}}+\left[8\hat{u}(\hat{s},\hat{m}_q)\left((1-\hat{m}_q^2)^2\right.\right.\\
&\left.-(1+\hat{m}_q^2)\hat{s}\right)+4(1-2\hat{m}_q^2+\hat{m}_q^4-\hat{s}-\hat{m}_q^2\hat{s})\,\hat{u}(\hat{s},\hat{m}_q)\,\hat{\lambda}_1\\
&+4\left(-5+30\hat{m}_q^4-40\hat{m}_q^6+15\hat{m}_q^8-\hat{s}+21\hat{m}_q^2\hat{s}+25\hat{m}_q^4\hat{s}-45\hat{m}_q^6\hat{s}+13\hat{s}^2\right.\\
&\left.\left.\left.+22\hat{m}_q^2\hat{s}^2+45\hat{m}_q^4\hat{s}^2-7\hat{s}^3-15\hat{m}_q^2\hat{s}^3\right)\frac{\hat{\lambda}_2}{\hat{u}(\hat{s},\hat{m}_q)}\right]Re(C_9^{\mathrm{eff}})\,C_7^{\mathrm{eff}}\right\}\,.
\end{aligned}
\tag{6}
$$

In the above expression, the branching ratio is normalized by $\mathcal{B}_{sl}$ for decays $B \to (X_c, X_u) l \nu_l$. We separately write a combination of the CKM factor $\frac{|V_{tq}^* V_{tb}|^2}{|V_{cb}|^2}$ due to top quark loop from the normalization factor $\mathcal{B}_0$. The normalization constant $\mathcal{B}_0$ is

$$\mathcal{B}_0 \equiv \mathcal{B}_{sl} \frac{3\,\alpha^2}{16\pi^2} \frac{1}{f(\hat{m}_c)\kappa(\hat{m}_c)}\,, \tag{7}$$

where $f \simeq 0.54$ is a phase space factor, and $\kappa(\hat{m}_c) \simeq 0.85$ includes $O(\alpha_s)$ QCD correction and $1/m_b^2$ corrections. The differential decay rate is not a simple parton model result. It contains non-perturbative $1/{m_b}^2$ power corrections and are denoted by the terms proportional to $\hat{\lambda}_1 \simeq -0.0087$ and $\hat{\lambda}_2 \simeq 0.0052$. Because of the smallness of these parameters, the correction is small compared with the leading parton model result for the region of $s << (m_b - m_q)^2$, $(q = d, s)$.

3 Numerical Results and Summary

From the present experimental value of $\Delta M_d = 0.474 \pm 0.031 (ps^{-1})$, and from the analysis on $B \to X_u l \bar{\nu}$, we obtain the limits on $(1-\rho)^2 + \eta^2$ and $\sqrt{\rho^2 + \eta^2}$:

$$\begin{aligned} (1-\rho)^2 + \eta^2 &= 0.85(1 \pm 0.15) \left\{ 0.2/({B_B}^{1/2} F_B(\mathrm{GeV})) \right\}^2 \\ &= [0.59 \pm 0.09, 1.3 \pm 0.20] \\ \sqrt{\rho^2 + \eta^2} &= 0.36 \pm 0.09, \end{aligned} \tag{8}$$

where we use the following values, $\eta_{QCD} = 0.55$, $m_t = 175(GeV)$ $A\lambda^2 = 0.041(\pm 0.003)$ and $\lambda = 0.2205, F_B = 0.2 \pm 0.04$. Within the limits, we show how $\frac{1}{\lambda^2}\frac{dR}{ds}$ depends on $(1-\rho)^2 + \eta^2$ and $\phi_1 = arg(-\frac{V_{cd} V_{cb}^*}{V_{td} V_{tb}^*})$.

In Fig.1, we show the ratio $\frac{1}{\lambda^2}\frac{dR(s)}{ds}$ for two sets of values for $(\phi_1, (1-\rho)^2 + \eta^2)$, one of which $(1-\rho)^2 + \eta^2$ is minimum and the other is maximum. The ratios for the CP conjugate process $\bar{B} \to \bar{X}_q l^+ l^-$ are also shown. To summarize the numerical results:

- The ratio at low invariant mass region $\simeq 1 (GeV^2)$ is very near to the value of $(1-\rho)^2 + \eta^2$, while on the peak of the resonance J/ψ, the ratio is almost 1. The value at low invariant mass region does not depend so much on whether the decaying particles are B or $\bar{B}$.

- For more details on this work, please see Ref. [4].



Figure 1: The ratio $\frac{1}{\lambda^2}\frac{dR(s)}{ds}$ versus s (GeV^2) with two different values of $(1-\rho)^2+\eta^2$. The solid curve corresponds to $(\phi_1,(1-\rho)^2+\eta^2)=(20,0.59)$, and the boxed curve corresponds to $(20,1.33)$. The ratio for CP conjugate process $\bar{B}\to\bar{X}_q l^+ l^-$ is denoted by the crossed curve for $(1-\rho)^2+\eta^2=1.33$, and by the dotted curve for $(1-\rho)^2+\eta^2=0.59$.

Acknowledgments

We would like to thank the organizer of BCONF97. The work of T.M. is supported by Monbusho Grant, No. 08044089. The work of C.S.K. is, in part, supported by BSRI-97-2425, and the work of A.I.S. was, in part, supported by Daiko Foundation.

References

1. C.S. Lim, T. Morozumi and A.I. Sanda, *Phys. Lett.* B **218**, 343 (1989).
2. A. Ali, G. Hiller, L. Handoko, T. Morozumi *Phys. Rev.* D **57**, 4105 (1997).
3. F. Krüger and L.M. Sehgal, *Phys. Rev.* D **55**, 2799 (1997).
4. C.S. Kim, T. Morozumi and A.I. Sanda, work in progress (1997).